

Protect your hearing & **your health**

The Hidden Toll of Occupational Noise Exposure

We know that exposure to too much noise damages hearing, but did you know that noise exposure is also associated with other conditions? The following are examples of how hazardous noise can affect your health.

Reduce noise exposure to protect your hearing AND your health!

Noise Induced Hearing Loss



Reduced hearing makes it difficult to hear conversations, important sounds, and to stay connected with your surroundings. Noise induced hearing loss is permanent, but completely preventable.

Mental Health

Anxiety, annoyance, stress and depression are linked with working in high noise.

Sleep Quality

Occupational noise exposure interferes with sleep patterns, which can increase the risk of heart disease, diabetes, and obesity.

Social Isolation

People may stop joining social activities because they have trouble hearing people speaking

Accidents/Injuries

Workplace noise is associated with an increased risk of accidents and injuries. This may be due to factors such as noise interfering with communication, worker distraction, stress, or fatigue.

Tinnitus


Ringing or buzzing in the ears, known as tinnitus, may be a sign of damage to the ears – similar to how sunburn is a warning sign of skin damage. Tinnitus can become a permanent condition.

Cardiovascular Disease

Working in noise is associated with higher risk of high blood pressure, elevated cholesterol, heart attacks and strokes.

Reproductive Health

If you are pregnant, or thinking of becoming pregnant, talk to your doctor about the risk of high noise exposure.



References:

- Themann CL, Masterson EA. Occupational noise exposure: A review of its effects, epidemiology, and impact with recommendations for reducing its burden. J. Acoust. Soc. 2019;146:3879-3905.
- Themann C, Suter AH, Stephenson MR. National Research Agenda for the Prevention of Occupational Hearing Loss – Part 1. Semin Hear. 2013;34(3):145-207.
- Rabinowitz, PM, Davies, HW, and Meinke, DK. AIHA Noise Manual 6th Edition Chapter 5 Auditory and Non-Auditory Health Effects of Noise. 2022.
- Kerns EA, Masterson EA, Themann CL, Calvert GM. Cardiovascular conditions, hearing difficulty, and occupational noise exposure within US industries and occupations. Am J Ind Med. 2018;61(6):477-491. doi:10.1002/ajim.22833
- Dementia prevention, intervention, and care: 2024 report of the Lancet standing Commission. Livingston, Gill et al. The Lancet, Volume 404, Issue 10452, 572 – 628
- Yoon JH, Roh J, Kim CN, Won JU. The risk of occupational injury increased according to severity of noise exposure after controlling for occupational environment status in Korea. Noise Health. 2016 Nov-Dec;18(85):355-361. doi: 10.4103/1463-1741.195811. PMID: 27991467; PMCID: PMC5227016.
- Girard SA, Leroux T, Verreault R, et al. Falls risk and hospitalization among retired workers with occupational noise-induced hearing loss. Can J Aging. 2014;33(1):84-91. doi:10.1017/S0714980813000664
- Yazdani-rad S, Khoshakhlagh AH, Al Sulaie S, Drake CL, Wickwire EM (2023). The effects of occupational noise on sleep: A systematic review. Sleep Medicine Reviews 72(2023). <https://doi.org/10.1016/j.smrv.2023.101846>



Through OSHA's Alliances with ASSP, CAOHC, and ISEA, the participants developed this infographic for informational purposes only. It does not necessarily reflect the official views of OSHA or the U.S. Department of Labor. August 2025.